

## GLYCOPEPTIDE DISULFIDE AND THIOESTER DERIVATIVES

### PRIORITY OF INVENTION

This application claims priority to United States Provisional Application  
Number 60/213,146, filed 22 June 2000, which application is incorporated herein by  
5 reference in its entirety.

### BACKGROUND OF THE INVENTION

#### Field of the Invention

This invention is directed to novel disulfide and thioester derivatives of  
glycopeptide antibiotics and related compounds. This invention is also directed to  
10 pharmaceutical compositions containing such glycopeptide derivatives, methods of  
using such glycopeptide derivatives as antibacterial agents, and processes and  
intermediates useful for preparing such glycopeptide derivatives.

#### Background

Glycopeptides are a well-known class of antibiotics produced by various  
15 microorganisms (*see Glycopeptide Antibiotics*, edited by R. Nagarajan, Marcel Dekker,  
Inc. New York (1994)). These complex multi-ring peptide compounds are very  
effective antibacterial agents against a majority of Gram-positive bacteria. Although  
potent antibacterial agents, the glycopeptides antibiotics are not used in the treatment of  
bacterial diseases as often as other classes of antibiotics, such as the semi-synthetic  
20 penicillins, cephalosporins and lincomycins, due to concerns regarding toxicity.

In recent years, however, bacterial resistance to many of the commonly-used  
antibiotics has developed (*see J. E. Geraci et al., Mayo Clin. Proc.* **1983**, 58, 88-91; and  
M. Foldes, *J. Antimicrob. Chemother.* **1983**, 11, 21-26). Since glycopeptide antibiotics

are often effective against these resistant strains of bacteria, glycopeptides such as vancomycin have become the drugs of last resort for treating infections caused by these organisms. Recently, however, resistance to vancomycin has appeared in various microorganisms, such as vancomycin-resistant enterococci (VRE), leading to increasing concerns about the ability to effectively treat bacterial infections in the future (see  
5 Hospital Infection Control Practices Advisory Committee, *Infection Control Hospital Epidemiology*, **1995**, *17*, 364-369; A. P. Johnson et al., *Clinical Microbiology Rev.*, **1990**, *3*, 280-291; G. M. Eliopoulos, *European J. Clinical Microbiol., Infection Disease*, **1993**, *12*, 409-412; and P. Courvalin, *Antimicrob. Agents Chemother*, **1990**, *34*, 2291-  
10 2296).

A number of derivatives of vancomycin and other glycopeptides are known in the art. For example, see U.S. Patent Nos. 4,639,433; 4,643,987; 4,497,802; 4,698,327; 5,591,714; 5,840,684; and 5,843,889. Other derivatives are disclosed in EP 0 802 199; EP 0 801 075; EP 0 667 353; WO 97/28812; WO 97/38702; WO 98/52589; WO  
15 98/52592; and in *J. Amer. Chem. Soc.*, **1996**, *118*, 13107-13108; *J. Amer. Chem. Soc.*, **1997**, *119*, 12041-12047; and *J. Amer. Chem. Soc.*, **1994**, *116*, 4573-4590.

Despite the above referenced disclosures, a need currently exists for novel glycopeptide derivatives having effective antibacterial activity and an improved mammalian safety profile. In particular, a need exists for glycopeptide derivatives  
20 which are effective against a wide spectrum of pathogenic microorganism, including vancomycin-resistant microorganisms, and which have reduced tissue accumulation and/or nephrotoxicity.

## SUMMARY OF THE INVENTION

The present invention provides novel disulfide and thioester glycopeptide  
25 derivatives having highly effective antibacterial activity and an improved mammalian safety profile. More specifically, the disulfide and thioester glycopeptide derivatives of

the invention unexpectedly exhibit reduced tissue accumulation and/or nephrotoxicity when administered to a mammal.

Accordingly, the invention provides a compound of the invention, which is a glycopeptide compound having at least one substituent of the formula:



wherein

each  $R^a$  is independently alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene, substituted alkynylene, cycloalkylene, substituted cycloalkylene, cycloalkenylene, substituted cycloalkenylene, arylene, heteroarylene, heterocyclene, 10 -C(O)-alkylene, substituted -C(O)-alkylene, -C(O)-alkenylene, substituted -C(O)-alkenylene, -C(O)-alkynylene, substituted -C(O)-alkynylene, -C(O)-cycloalkylene, substituted -C(O)-cycloalkylene, -C(O)-cycloalkenylene, substituted -C(O)-cycloalkenylene, -C(O)-arylene, -C(O)-heteroarylene, or -C(O)-heterocyclene;

15 each  $R^b$  is independently a covalent bond, alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene, substituted alkynylene, cycloalkylene, substituted cycloalkylene, cycloalkenylene, or substituted cycloalkenylene; provided  $R^b$  is not a covalent bond when Z is hydrogen;

each Y is independently selected from the group consisting of oxygen, sulfur, 20 -S-S-, -S-C(=O)-, -C(=O)-S-, -NR<sup>c</sup>-, -S(O)-, -SO<sub>2</sub>-, -NR<sup>c</sup>C(O)-, -OSO<sub>2</sub>-, -OC(O)-, -NR<sup>c</sup>SO<sub>2</sub>-, -C(O)NR<sup>c</sup>-, -C(O)O-, -SO<sub>2</sub>NR<sup>c</sup>-, -SO<sub>2</sub>O-, -P(O)(OR<sup>c</sup>)O-, -P(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OP(O)(OR<sup>c</sup>)O-, -OP(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OC(O)O-, -NR<sup>c</sup>C(O)O-, -NR<sup>c</sup>C(O)NR<sup>c</sup>-, -OC(O)NR<sup>c</sup>-, C(=O), and -NR<sup>c</sup>SO<sub>2</sub>NR<sup>c</sup>-;

25 each Z is independently selected from hydrogen, aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic;

each R<sup>c</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic and -C(O)R<sup>d</sup>;

- 5            each R<sup>d</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

*x* is 1 or 2;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;


- 10           provided that at least one Y is -S-S-, -S-C(=O)-, or -C(=O)-S-.

Preferably, each R<sup>a</sup> is independently selected from the group consisting of alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene.

- 15           Preferably, each R<sup>b</sup> is independently selected from the group consisting of a covalent bond, alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene, provided R<sup>b</sup> is not a covalent bond when Z is hydrogen.

Preferred compounds of the invention exclude glycopeptides substituted at the carboxy terminus with a substituent that comprises more than one carboxy group.

- 20           A preferred compound of the invention is a glycopeptide of formula I:



(I)

wherein:

R<sup>1</sup> is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic, -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>; or a saccharide group optionally substituted with -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>;

R<sup>2</sup> is hydrogen or a saccharide group optionally substituted with -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>, R<sup>f</sup>, -C(O)R<sup>f</sup>, or -C(O)-R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>;

R<sup>3</sup> is -OR<sup>c</sup>, -NR<sup>c</sup>R<sup>c</sup>, -O-R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>, -NR<sup>c</sup>-R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>, -NR<sup>c</sup>R<sup>e</sup>, or -O-R<sup>e</sup>;

R<sup>4</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>, -C(O)R<sup>d</sup> and

a saccharide group optionally substituted with  $-R^a-Y-R^b-(Z)_x$ ,  $R^f$ ,  $-C(O)R^f$ , or  $-C(O)-R^a-Y-R^b-(Z)_x$ ;

$R^5$  is selected from the group consisting of hydrogen, halo,  $-CH(R^c)-NR^cR^c$ ,  $-CH(R^c)-NR^cR^e$ ,  $-CH(R^c)-NR^c-R^a-Y-R^b-(Z)_x$ ,  $-CH(R^c)-R^x$ , and

5  $-CH(R^c)-NR^c-R^a-C(=O)-R^x$ ;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  $-R^a-Y-R^b-(Z)_x$ ,  $-C(O)R^d$  and a saccharide group optionally substituted with  $-NR^c-R^a-Y-R^b-(Z)_x$ , or  $R^5$  and  $R^6$  can be joined, together with the atoms to which they are attached, form a heterocyclic ring optionally substituted with  $-NR^c-R^a-Y-R^b-(Z)_x$ ;

10

$R^7$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  $-R^a-Y-R^b-(Z)_x$  and  $-C(O)R^d$ ;

$R^8$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15

$R^9$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

$R^{10}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic; or  $R^8$  and  $R^{10}$  are joined to form  $-Ar^1-O-Ar^2-$ , where  $Ar^1$  and  $Ar^2$  are independently arylene or heteroarylene;

20

$R^{11}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic, or

25

R<sup>10</sup> and R<sup>11</sup> are joined, together with the carbon and nitrogen atoms to which they are attached, to form a heterocyclic ring;

R<sup>12</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic,  
5 -C(O)R<sup>d</sup>, -C(NH)R<sup>d</sup>, -C(O)NR<sup>c</sup>R<sup>c</sup>, -C(O)OR<sup>d</sup>, -C(NH)NR<sup>c</sup>R<sup>c</sup> and -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub>, or  
R<sup>11</sup> and R<sup>12</sup> are joined, together with the nitrogen atom to which they are attached, to form a heterocyclic ring;

R<sup>13</sup> is selected from the group consisting of hydrogen or -OR<sup>14</sup>;

10 R<sup>14</sup> is selected from hydrogen, -C(O)R<sup>d</sup> and a saccharide group;

each R<sup>a</sup> is independently selected from the group consisting of alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene;

each R<sup>b</sup> is independently selected from the group consisting of a covalent bond, alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene, provided R<sup>b</sup> is not a covalent bond when Z is hydrogen;

each R<sup>c</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl,  
20 heterocyclic and -C(O)R<sup>d</sup>;

each R<sup>d</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

R<sup>e</sup> is a saccharide group;

25 each R<sup>f</sup> is independently alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, or heterocyclic;

R<sup>x</sup> is a nitrogen-linked amino saccharide or a nitrogen-linked heterocycle;

X<sup>1</sup>, X<sup>2</sup> and X<sup>3</sup> are independently selected from hydrogen or chloro;

each Y is independently selected from the group consisting of oxygen, sulfur,

- 5    -S-S-, -S-C(=O)-, -C(=O)-S-, -NR<sup>c</sup>-, -S(O)-, -SO<sub>2</sub>-, -NR<sup>c</sup>C(O)-, -OSO<sub>2</sub>-,  
-OC(O)-, -NR<sup>c</sup>SO<sub>2</sub>-, -C(O)NR<sup>c</sup>-, -C(O)O-, -SO<sub>2</sub>NR<sup>c</sup>-, -SO<sub>2</sub>O-, -P(O)(OR<sup>c</sup>)O-,  
-P(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OP(O)(OR<sup>c</sup>)O-, -OP(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OC(O)O-, -NR<sup>c</sup>C(O)O-,  
-NR<sup>c</sup>C(O)NR<sup>c</sup>-, -OC(O)NR<sup>c</sup>-, C(=O), and -NR<sup>c</sup>SO<sub>2</sub>NR<sup>c</sup>-;

each Z is independently selected from hydrogen, aryl, cycloalkyl, cycloalkenyl,  
heteroaryl and heterocyclic;

- 10    *n* is 0, 1 or 2; and

*x* is 1 or 2;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;

wherein the glycopeptide is substituted with one or more groups wherein Y is

-S-S-, -S-C(=O)-, or -C(=O)-S-;

- 15    provided R<sup>3</sup> is not a substituent that comprises more than one carboxy group; and  
provided R<sup>3</sup> is not a substituent that comprises one or more saccharide groups  
and a carboxy group; and

provided R<sup>1</sup> is not an amino saccharide wherein the saccharide-amine is  
substituted with a substituent that comprises two or more hydroxy groups.

- 20    Preferably, R<sup>1</sup> is an amino saccharide wherein the saccharide-amine is  
substituted with: -CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>11</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>10</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>;  
25    -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>3</sub>-CH=CH-(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> (*trans*); -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>-S(O)-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>6</sub>Ph; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>Ph;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>Ph; -CH<sub>2</sub>CH<sub>2</sub>-NH-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph;



$-\text{CH}_2\text{CH}_2-\text{NH}-\text{CH}_2-4-[4-(\text{CH}_3)_2\text{CHCH}_2-]-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2-\text{NH}-\text{CH}_2-4-(4-\text{CF}_3-\text{Ph})-\text{Ph}$ ;  
 $-\text{CH}_2\text{CH}_2-\text{S}-\text{CH}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2-\text{S}(\text{O})-\text{CH}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  
 $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{S}-\text{CH}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{S}(\text{O})-\text{CH}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  
 $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{S}-\text{CH}_2-4-[3,4-\text{di}-\text{Cl}-\text{PhCH}_2\text{O}-]-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2-\text{NHSO}_2-\text{CH}_2-4-[4-(4-$   
5  $\text{Ph})-\text{Ph}]-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{NHSO}_2-\text{CH}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  
 $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{NHSO}_2-\text{CH}_2-4-(\text{Ph}-\text{C}\equiv\text{C})-\text{Ph}$ ;  $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{NHSO}_2-4-(4-\text{Cl}-\text{Ph})-\text{Ph}$ ;  
or  $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{NHSO}_2-4-(\text{naphth-2-yl})-\text{Ph}$ . More preferably, the alkylated  
glycopeptide product is also compound of formula I wherein  $\text{R}^1$  is an amino saccharide  
wherein the saccharide-amine is substituted with 4-(4-chlorophenyl)benzyl or 4-(4-  
10 chlorobenzyloxy)benzyl.


More preferably,  $\text{R}^1$  is an amino containing saccharide group substituted on the  
 amine with a substituent that comprises one or more (e.g. 1, 2, or 3) disulfide ( $-\text{S}-\text{S}-$ ) or  
 thioester ( $-\text{S}-\text{C}(=\text{O})-$ ) bonds.

More preferably,  $\text{R}^1$  is an amino containing saccharide group substituted on the  
 15 amine with a group of formula  $-\text{R}^a-\text{W}-\text{R}^b$  wherein: W is  $-\text{S}-\text{S}-$  or  $-\text{S}-\text{C}(=\text{O})-$  and  $\text{R}^h$  is  
 alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  
 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl,  
 heteroaryl, or heterocyclic.

More preferably,  $\text{R}^a$  is alkylene, substituted alkylene, alkenylene, substituted  
 20 alkenylene, alkynylene, substituted alkynylene,  $-\text{C}(\text{O})$ -alkylene, substituted  
 $-\text{C}(\text{O})$ -alkylene,  $-\text{C}(\text{O})$ -alkenylene, substituted  $-\text{C}(\text{O})$ -alkenylene,  $-\text{C}(\text{O})$ -alkynylene, or  
 substituted  $-\text{C}(\text{O})$ -alkynylene.

More preferably,  $\text{R}^b$  is alkyl, substituted alkyl, alkenyl, substituted alkenyl,  
 alkynyl, or substituted alkynyl.

25 For example,  $\text{R}^1$  is preferably a saccharide group of the formula (III):



wherein  $R^{15}$  is  $-R^a-W-R^h$ ; and  $R^{16}$  is hydrogen or methyl.

Preferably,  $R^2$  is hydrogen.

In a preferred embodiment,  $R^3$  is  $-R^a-W-R^h$  where  $R^a$ ,  $R^h$ , and  $W$  are as defined herein. In another preferred embodiment,  $R^3$  is  $-OR^c$  or  $-NR^cR^c$ ; more preferably  $R^3$  is  $-OH$ .

- 5 Preferably,  $R^3$  is  $-OH$ ;  $-NH-(CH_2)_3-N(CH_3)_2$ ;  $N$ -(D-glucosamine);  $-NHCH(CO_2CH_3)CH_2CO_2CH_3$ ;  $-NH(CH_2)_3$ -(morpholin-4-yl);  $-NH(CH_2)_3-NH(CH_2)_2CH_3$ ;  $-NH(CH_2)$ -piperidin-1-yl;  $-NH(CH_2)_4NHC(N)NH_2$ ;  $-NH(CH_2)_2-N^+(CH_3)_3$ ;  $-NHCH(COOH)(CH_2)_3NHC(N)NH_2$ ;  $-NH-[(CH_2)_3-NH-]_3-H$ ;  $-N[(CH_2)_3N(CH_3)_2]_2$ ;
- 10  $-NH(CH_2)_3$ -imidazol-1-yl;  $-NHCH_2$ -4-pyridyl;  $-NH(CH_2)_3CH_3$ ;  $-NH(CH_2)_2OH$ ;  $-NH(CH_2)_5OH$ ;  $-NH(CH_2)_2OCH_3$ ;  $-NHCH_2$ -tetrahydrofuran-2-yl;  $-N[(CH_2)_2OH]_2$ ;  $-NH(CH_2)_2N[(CH_2)_2OH]_2$ ;  $-NHCH_2COOH$ ;  $-NHCH(COOH)CH_2OH$ ;  $-NH(CH_2)_2COOH$ ;  $N$ -(glucamine);  $-NH(CH_2)_2COOH$ ;  $-NH(CH_2)_3SO_3H$ ;  $-NHCH(COOH)(CH_2)_2NH_2$ ;  $-NHCH(COOH)(CH_2)_3NH_2$ ;
- 15  $-NHCH(COOH)CH_2CO_2(CH_2)_3-N^+(CH_3)_3$ ;  $-NHCH(COOH)CH_2CO_2(CH_2)_2C(O)-N(CH_3)_2$ ;  $-NHCH(COOH)CH_2CO_2(CH_2)_3$ -morpholin-4-yl;  $-NHCH(COOH)CH_2CO_2(CH_2)_2OC(O)C(CH_3)_3$ ;  $-NHCH(CH_2COOH)CO_2(CH_2)_3-$

$N^+(CH_3)_3$ ;  $-NHCH(CH_2COOH)CO_2(CH_2)_2C(O)N(CH_3)_2$ ;  
 $-NHCH(CH_2COOH)CO_2(CH_2)_3$ -morpholin-4-yl;  
 $-NHCH(CH_2COOH)CO_2(CH_2)_2OC(O)C(CH_3)_3$ ;  $-NHCH(COOH)CH_2CO_2CH_3$ ;  
 $-NHCH(CH_2COOH)CO_2(CH_2)_2N(CH_3)_2$ ;  $-NHCH(COOH)CH_2CO_2CH_2C(O)N(CH_3)_2$ ;  
5  $-NHCH(CH_2COOH)CO_2CH_2C(O)N(CH_3)_2$ ;  $-NHCH(CH_2COOH)CO_2CH_3$ ;  
 $-NH(CH_2)_3N(CH_3)_2$ ;  $-NHCH_2CH_2CO_2CH_3$ ;  $-NHCH[CH_2CO_2CH_2C(O)N(CH_3)_2]$ -  
 $CO_2CH_2-C(O)-N(CH_3)_2$ ;  $-NHCH_2CO_2CH_3$ ;  $-N$ -(methyl 3-amino-3-  
deoxyaminopyranoside);  $-N$ -(methyl 3-amino-2,3,6-trideoxyhexopyranoside);  $-N$ -(2-  
amino-2-deoxy-6-(dihydrogenphosphate)-glucopyranose);  $-N$ -(2-amino-2-deoxygluconic  
10 acid);  $-NH(CH_2)_4COOH$ ;  $-N$ -( $N$ -CH<sub>3</sub>-D-glucamine);  $-NH(CH_2)_6COOH$ ;  $-O$ (D-glucose);  
 $-NH(CH_2)_3OC(O)CH(NH_2)CH_3$ ;  $-NH(CH_2)_4CH(C(O)-2-HOOC-pyrrolidin-1-$   
yl) $NHCH(COOH)-CH_2CH_2Ph$  (S,S isomer);  $-NH-CH_2CH_2-NH-(CH_2)_9CH_3$ ;  
 $-NH(CH_2)C(O)CH_2C(O)N(CH_3)_2$ ;

Preferably,  $R^4$ ,  $R^6$  and  $R^7$  are each independently selected from hydrogen or  
15  $-C(O)R^d$ . More preferably,  $R^4$ ,  $R^6$  and  $R^7$  are each hydrogen.

In a preferred embodiment,  $R^5$  is  $-R^a-W-R^h$  where  $R^a$ ,  $R^h$ , and  $W$  are as defined  
 herein. In another preferred embodiment,  $R^5$  is hydrogen,  $-CH_2-NHR^c$ ,  $-CH_2-NR^cR^e$  or  
 $-CH_2-NH-R^a-Y-R^b-(Z)_x$ .  $R^5$  can also preferably be hydrogen;  $-CH_2-N$ -( $N$ -CH<sub>3</sub>-D-  
glucamine);  $-CH_2-NH-CH_2CH_2-NH-(CH_2)_9CH_3$ ;  $-CH_2-NH-CH_2CH_2-NHC(O)-$   
20  $(CH_2)_3COOH$ ;  $-CH_2-NH-(CH_2)_9CH_3$ ;  $-CH_2-NH-CH_2CH_2-COOH$ ;  $-CH_2-NH-$   
 $(CH_2)_5COOH$ ;  $-CH_2$ -(morpholin-4-yl);  $-CH_2-NH-CH_2CH_2-O-CH_2CH_2OH$ ;  $-CH_2-NH-$   
 $CH_2CH(OH)-CH_2OH$ ;  $-CH_2-N[CH_2CH_2OH]_2$ ;  $-CH_2-NH-(CH_2)_3-N(CH_3)_2$ ;  $-CH_2-$   
 $N[(CH_2)_3-N(CH_3)_2]_2$ ;  $-CH_2-NH-(CH_2)_3$ -(imidazol-1-yl);  $-CH_2-NH-(CH_2)_3$ -(morpholin-4-  
yl);  $-CH_2-NH-(CH_2)_4-NHC(NH)NH_2$ ;  $-CH_2-N$ -(2-amino-2-deoxygluconic acid);  $-CH_2-$   
25  $NH-CH_2CH_2-NH-(CH_2)_{11}CH_3$ ;  $-CH_2-NH-CH(COOH)CH_2COOH$ ;  $-CH_2-NH-CH_2CH_2-$   
 $NHSO_2-(CH_2)_7CH_3$ ;  $-CH_2-NH-CH_2CH_2-NHSO_2-(CH_2)_8CH_3$ ;  $-CH_2-NH-CH_2CH_2-$   
 $NHSO_2-(CH_2)_9CH_3$ ;  $-CH_2-NH-CH_2CH_2-NHSO_2-(CH_2)_{11}CH_3$ ;  $-CH_2-NH-CH_2CH_2-NH-$

(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>2</sub>CH<sub>2</sub>OH; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>C(O)-N-(D-glucosamine); -CH<sub>2</sub>-NH-(6-oxo-[1,3]oxazinan-3-yl); -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>11</sub>CH<sub>3</sub>; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>6</sub>Ph; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>Ph; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>10</sub>Ph; -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>2</sub>-(4-(4-CF<sub>3</sub>-Ph)Ph); -CH<sub>2</sub>-NH-CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>11</sub>CH<sub>3</sub>; or -CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>5</sub>-COOH.

Preferably, R<sup>8</sup> is -CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>COOH, benzyl, 4-hydroxyphenyl or 3-chloro-4-hydroxyphenyl.

Preferably, R<sup>9</sup> is hydrogen or alkyl.

10 Preferably, R<sup>10</sup> is alkyl or substituted alkyl. More preferably, R<sup>10</sup> is the side-chain of a naturally occurring amino acid, such as isobutyl.

Preferably, R<sup>11</sup> is hydrogen or alkyl.

15 Preferably, R<sup>12</sup> is hydrogen, alkyl, substituted alkyl or -C(O)R<sup>d</sup>. R<sup>12</sup> can also preferably be hydrogen; -CH<sub>2</sub>COOH; -CH<sub>2</sub>-[CH(OH)]<sub>5</sub>CH<sub>2</sub>OH; -CH<sub>2</sub>CH(OH)CH<sub>2</sub>OH; -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>; -CH<sub>2</sub>C(O)OCH<sub>2</sub>CH<sub>3</sub>; -CH<sub>2</sub>-(2-pyridyl); -CH<sub>2</sub>-[CH(OH)]<sub>4</sub>COOH; -CH<sub>2</sub>-(3-carboxyphenyl); (*R*)-C(O)CH(NH<sub>2</sub>)(CH<sub>2</sub>)<sub>4</sub>NH<sub>2</sub>; -C(O)Ph; -C(O)CH<sub>2</sub>NHC(O)CH<sub>3</sub>; *E*-CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>3</sub>CH=CH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>; or -C(O)CH<sub>3</sub>.


Preferably, X<sup>1</sup> and X<sup>2</sup> are each chloro.

Preferably, X<sup>3</sup> is hydrogen.

20 Preferably, each Y is independently selected from the group consisting of oxygen, sulfur, -S-S-, -S-C(=O)-, -C(=O)-S-, -NR<sup>c</sup>-, -S(O)-, -SO<sub>2</sub>-, -NR<sup>c</sup>C(O)-, -OSO<sub>2</sub>-, -OC(O)-, -NR<sup>c</sup>SO<sub>2</sub>-, -C(O)NR<sup>c</sup>-, -C(O)O-, -SO<sub>2</sub>NR<sup>c</sup>-, -SO<sub>2</sub>O-, -P(O)(OR<sup>c</sup>)O-, -P(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OP(O)(OR<sup>c</sup>)O-, -OP(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OC(O)O-, -NR<sup>c</sup>C(O)O-, -NR<sup>c</sup>C(O)NR<sup>c</sup>-, -OC(O)NR<sup>c</sup>-, and -NR<sup>c</sup>SO<sub>2</sub>NR<sup>c</sup>-;

25 Preferably, *n* is 0 or 1, and more preferably, *n* is 1.

A preferred compound of the invention is a glycopeptide of formula II:



(II)

wherein:

$R^{19}$  is hydrogen;

$R^{20}$  is  $-R^a-W-R^h$

$R^a$  is independently alkylene, substituted alkylene, alkenylene, substituted  
 5 alkenylene, alkynylene, substituted alkynylene, cycloalkylene, substituted cycloalkylene,  
 cycloalkenylene, substituted cycloalkenylene, arylene, heteroarylene, heterocyclene,  
 -C(O)-alkylene, substituted -C(O)-alkylene, -C(O)-alkenylene, substituted  
 -C(O)-alkenylene, -C(O)-alkynylene, substituted -C(O)-alkynylene,  
 -C(O)-cycloalkylene, substituted -C(O)-cycloalkylene, -C(O)-cycloalkenylene,  
 10 substituted -C(O)-cycloalkenylene, -C(O)-arylene, -C(O)-heteroarylene, or  
 -C(O)-heterocyclene;

R<sup>h</sup> is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, or heterocyclic;

W is -S-S-, -C(=O)S-, or -S-C(=O)- and

- 5 R<sup>3</sup>, and R<sup>5</sup> have any of the values or preferred values described herein;  
or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;  
provided R<sup>3</sup> is not a substituent that comprises more than one carboxy group;  
and

- 10 provided R<sup>3</sup> is not a substituent that comprises one or more saccharide groups  
and a carboxy group; and

provided R<sup>20</sup> is not a substituent that comprises two or more hydroxy groups.

- More preferably, for a compound of formula II, R<sup>a</sup> is alkylene, substituted  
alkylene, alkenylene, substituted alkenylene, alkynylene, substituted alkynylene,  
-C(O)-alkylene, substituted -C(O)-alkylene, -C(O)-alkenylene, substituted  
15 -C(O)-alkenylene, -C(O)-alkynylene, or substituted -C(O)-alkynylene; and R<sup>h</sup> is alkyl,  
substituted alkyl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl.

- A preferred value for R<sup>15</sup>, R<sup>20</sup>, or -R<sup>a</sup>-Y-R<sup>b</sup>-(Z)<sub>x</sub> is -CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>11</sub>CH<sub>3</sub>;  
20 -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>10</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>3</sub>-  
CH=CH-(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> (*trans*); -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>;  
-CH<sub>2</sub>CH<sub>2</sub>-S(O)-(CH<sub>2</sub>)<sub>9</sub>CH<sub>3</sub>; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>6</sub>Ph; -CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>Ph;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-(CH<sub>2</sub>)<sub>8</sub>Ph; -CH<sub>2</sub>CH<sub>2</sub>-NH-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph;  
25 -CH<sub>2</sub>CH<sub>2</sub>-NH-CH<sub>2</sub>-4-[4-(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>-]-Ph; -CH<sub>2</sub>CH<sub>2</sub>-NH-CH<sub>2</sub>-4-(4-CF<sub>3</sub>-Ph)-Ph;  
-CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph; -CH<sub>2</sub>CH<sub>2</sub>-S(O)-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S(O)-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph;

-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>2</sub>-4-[3,4-di-Cl-PhCH<sub>2</sub>O-)-Ph; -CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-CH<sub>2</sub>-4-[4-(4-Ph)-Ph]-Ph; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-CH<sub>2</sub>-4-(4-Cl-Ph)-Ph;  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-CH<sub>2</sub>-4-(Ph-C≡C-)-Ph; -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-4-(4-Cl-Ph)-Ph;  
or -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-NHSO<sub>2</sub>-4-(naphth-2-yl)-Ph. Another preferred value for R<sup>15</sup> or R<sup>20</sup>

5 4-(4-chlorophenyl)benzyl or 4-(4-chlorobenzoyloxy)benzyl

The invention also provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of the invention. In one preferred embodiment, the pharmaceutically acceptable carrier comprises an aqueous cyclodextrin solution. Preferably, the  
10 cyclodextrin is hydroxypropyl-β-cyclodextrin or sulfobutyl ether β-cyclodextrin. More preferably, the cyclodextrin is hydroxypropyl-β-cyclodextrin.

The compounds of the invention are highly effective antibacterial agents. Accordingly, the invention also provides a method of treating a mammal having a bacterial disease, comprising administering to the mammal a therapeutically effective  
15 amount of a compound of the invention. The invention also provides a method of treating a mammal having a bacterial disease, comprising administering to the mammal a therapeutically effective amount of a pharmaceutical composition of the invention.

The invention also provides processes and intermediates useful for preparing compounds of the invention, which processes and intermediates are described further  
20 herein.

The invention also provides a compound of the invention as described herein for use in medical therapy, as well as the use of a compound of the invention in the manufacture of a formulation or medicament for treating a bacterial disease in a mammal.

25 Preferred compounds of the invention are the compounds of formula II shown in Table I below wherein R<sup>19</sup> is hydrogen.

Table I: Preferred Compounds of formula II

Compound	R <sup>3</sup>	R <sup>5</sup>	R <sup>20</sup>
<b>1</b>	OH	H	-(CH <sub>2</sub> ) <sub>2</sub> S-S(CH <sub>2</sub> ) <sub>7</sub> CH <sub>3</sub>
<b>2</b>	OH	H	-(CH <sub>2</sub> ) <sub>3</sub> S-S(CH <sub>2</sub> ) <sub>7</sub> CH <sub>3</sub>
<b>3</b>	OH	H	-(CH <sub>2</sub> ) <sub>2</sub> S-S(CH <sub>2</sub> ) <sub>8</sub> CH <sub>3</sub>
<b>4</b>	OH	H	-(CH <sub>2</sub> ) <sub>3</sub> S-S(CH <sub>2</sub> ) <sub>8</sub> CH <sub>3</sub>

Another preferred group of compounds of the invention are disulfide and thioester derivatives of the glycopeptide antibiotic A82846B (also known as chloroorienticin A or LY264826). See for example R. Nagarajan et al., *J. Org. Chem.*, **1988**, *54*, 983-986; and N. Tsuji et al., *J. Antibiot.*, **1988**, *41*, 819-822. The structure of this glycopeptide is similar to vancomycin, except A82846B contains an additional amino sugar (i.e. 4-epi-vancosamine attached at the R<sup>2</sup> position in formula I.) and further contains 4-epi-vancosamine in place of vancosamine in the disaccharide moiety attached at the R<sup>1</sup> position in formula I. For example, a preferred group of compounds are derivatives of A82846B that are substituted at the 4-epi-vancosamine nitrogen with a substituent that comprises one or more disulfide and thioester groups; or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof. Another preferred group of compounds of the invention are derivatives of A82846B wherein the 4-epi-vancosamine has been replaced with a group selected from the values defined herein for R<sup>1</sup> in a compound of formula I. The compounds of the invention that are disulfide and thioester derivatives of A82846B can readily be prepared using the procedures described herein. Other preferred compounds of the invention are derivatives of A82846B wherein the 4-epi-vancosamine nitrogen is substituted with a 4-(4-chlorophenyl)benzyl group or a 4-(4-chlorobenzoyloxy)benzyl group.

Representative compounds of the invention have been found to unexpectedly



exhibit reduced tissue accumulation and/or nephrotoxicity when administered to a mammal. While not wishing to be bound by theory, it is believed that the disulfide or the thioester linkage functions as a metabolic sight that can be degraded *in vivo*, thereby facilitating excretion from the mammal after administration. The unexpected increase  
5 in excretion of the compounds of the invention may be responsible for the reduced tissue accumulation and/or reduced nephrotoxicity observed for these compounds relative to the corresponding compounds that lack the disulfide or thioester functionality.

Additionally, it is possible that the thiol metabolites resulting from the metabolic  
10 degradation of the compounds of the invention can undergo dimerization to form glycopeptide dimers that possess antibacterial activity. Accordingly, the invention also provides dimers of the formula  $G-R^a-B-R^a-G$ , wherein each G is independently a glycopeptide of formula I, bonded at the nitrogen of an  $R^1$  saccharide; each  $R^a$  independently has one of the values defined herein for  $R^a$  in a compound of formula I;  
15 and B is -S-S-.

### DETAILED DESCRIPTION OF THE INVENTION

This invention relates to novel compounds of the invention, which are disulfide or thioester derivatives of glycopeptide antibiotics, as well as to compositions comprising such compounds and to therapeutic methods comprising the administration  
20 of such compounds. When describing the compounds, compositions and methods of the invention, the following terms have the following meanings, unless otherwise indicated.

#### Definitions

The term "alkyl" refers to a monoradical branched or unbranched saturated hydrocarbon chain preferably having from 1 to 40 carbon atoms, more preferably 1 to 10  
25 carbon atoms, and even more preferably 1 to 6 carbon atoms. This term is exemplified

by groups such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *n*-hexyl, *n*-decyl, tetradecyl, and the like.

The term "substituted alkyl" refers to an alkyl group as defined above, having from 1 to 8 substituents, preferably 1 to 5 substituents, and more preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, guanido, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, -SO<sub>3</sub>H, and -SO<sub>2</sub>-heteroaryl.

The term "alkylene" refers to a diradical of a branched or unbranched saturated hydrocarbon chain, preferably having from 1 to 40 carbon atoms, preferably 1-10 carbon atoms, more preferably 1-6 carbon atoms. This term is exemplified by groups such as methylene (-CH<sub>2</sub>-), ethylene (-CH<sub>2</sub>CH<sub>2</sub>-), the propylene isomers (e.g., -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- and -CH(CH<sub>3</sub>)CH<sub>2</sub>-) and the like.

The term "substituted alkylene" refers to an alkylene group, as defined above, having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-

heteroaryl. Additionally, such substituted alkylene groups include those where 2 substituents on the alkylene group are fused to form one or more cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heterocyclic or heteroaryl groups fused to the alkylene group. Preferably such fused groups contain from 1 to 3 fused ring structures. Additionally, the term substituted alkylene includes alkylene groups in which from 1 to 5 of the alkylene carbon atoms are replaced with oxygen, sulfur or -NR- where R is hydrogen or alkyl. Examples of substituted alkenes are chloromethylene (-CH(Cl)-), aminoethylene (-CH(NH<sub>2</sub>)CH<sub>2</sub>-), 2-carboxypropylene isomers (-CH<sub>2</sub>CH(CO<sub>2</sub>H)CH<sub>2</sub>-), ethoxyethyl (-CH<sub>2</sub>CH<sub>2</sub> O-CH<sub>2</sub>CH<sub>2</sub>-) and the like.

10 The term "alkaryl" refers to the groups -alkylene-aryl and -substituted alkylene-aryl where alkylene, substituted alkylene and aryl are defined herein. Such alkaryl groups are exemplified by benzyl, phenethyl and the like.

The term "alkoxy" refers to the groups alkyl-O-, alkenyl-O-, cycloalkyl-O-, cycloalkenyl-O-, and alkynyl-O-, where alkyl, alkenyl, cycloalkyl, cycloalkenyl, and alkynyl are as defined herein. Preferred alkoxy groups are alkyl-O- and include, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

20 The term "substituted alkoxy" refers to the groups substituted alkyl-O-, substituted alkenyl-O-, substituted cycloalkyl-O-, substituted cycloalkenyl-O-, and substituted alkynyl-O- where substituted alkyl, substituted alkenyl, substituted cycloalkyl, substituted cycloalkenyl and substituted alkynyl are as defined herein.

The term "alkylalkoxy" refers to the groups -alkylene-O-alkyl, alkylene-O-substituted alkyl, substituted alkylene-O-alkyl and substituted alkylene-O-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein. Preferred alkylalkoxy groups are alkylene-O-alkyl and include, by way of example, methylenemethoxy (-CH<sub>2</sub>OCH<sub>3</sub>), ethylenemethoxy (-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>), *n*-propylene-*iso*-propoxy (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH(CH<sub>3</sub>)<sub>2</sub>), methylene-*t*-butoxy (-CH<sub>2</sub>-O-

C(CH<sub>3</sub>)<sub>3</sub>) and the like.

The term "alkylthioalkoxy" refers to the group -alkylene-S-alkyl, alkylene-S-substituted alkyl, substituted alkylene-S-alkyl and substituted alkylene-S-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein. Preferred alkylthioalkoxy groups are alkylene-S-alkyl and include, by way of example, methylenethiomethoxy (-CH<sub>2</sub>SCH<sub>3</sub>), ethylenethiomethoxy (-CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>), *n*-propylene-*iso*-thiopropoxy (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>SCH(CH<sub>3</sub>)<sub>2</sub>), methylene-*t*-thiobutoxy (-CH<sub>2</sub>SC(CH<sub>3</sub>)<sub>3</sub>) and the like.

The term "alkenyl" refers to a monoradical of a branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of vinyl unsaturation. Preferred alkenyl groups include ethenyl (-CH=CH<sub>2</sub>), *n*-propenyl (-CH<sub>2</sub>CH=CH<sub>2</sub>), *iso*-propenyl (-C(CH<sub>3</sub>)=CH<sub>2</sub>), and the like.

The term "substituted alkenyl" refers to an alkenyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-heteroaryl.

The term "alkenylene" refers to a diradical of a branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and

having at least 1 and preferably from 1-6 sites of vinyl unsaturation. This term is exemplified by groups such as ethenylene ( $-\text{CH}=\text{CH}-$ ), the propenylene isomers (e.g.,  $-\text{CH}_2\text{CH}=\text{CH}-$  and  $-\text{C}(\text{CH}_3)=\text{CH}-$ ) and the like.

The term "substituted alkenylene" refers to an alkenylene group as defined above  
5 having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol,  
10 thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro,  $-\text{SO}-$ alkyl,  $-\text{SO}-$ substituted alkyl,  $-\text{SO}-$ aryl,  $-\text{SO}-$ heteroaryl,  $-\text{SO}_2-$ alkyl,  $-\text{SO}_2-$ substituted alkyl,  $-\text{SO}_2-$ aryl and  $-\text{SO}_2-$ heteroaryl. Additionally, such substituted alkenylene groups include those where 2 substituents on the alkenylene group are fused to form one or more cycloalkyl,  
15 substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heterocyclic or heteroaryl groups fused to the alkenylene group.

The term "alkynyl" refers to a monoradical of an unsaturated hydrocarbon preferably having from 2 to 40 carbon atoms, more preferably 2 to 20 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6  
20 sites of acetylene (triple bond) unsaturation. Preferred alkynyl groups include ethynyl ( $-\text{C}\equiv\text{CH}$ ), propargyl ( $-\text{CH}_2\text{C}\equiv\text{CH}$ ) and the like.

The term "substituted alkynyl" refers to an alkynyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl,  
25 cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol,

thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-heteroaryl.

5           The term "alkynylene" refers to a diradical of an unsaturated hydrocarbon preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of acetylene (triple bond) unsaturation. Preferred alkynylene groups include ethynylene (-C≡C-), propargylene (-CH<sub>2</sub>C≡C-) and the like.

10           The term "substituted alkynylene" refers to an alkynylene group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, 15 thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-heteroaryl

20           The term "acyl" refers to the groups HC(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, cycloalkenyl-C(O)-, substituted cycloalkenyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic are as defined herein.

25           The term "acylamino" or "aminocarbonyl" refers to the group -C(O)NRR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic or where both R groups are joined to form a heterocyclic group (e.g., morpholino)

wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

The term "aminoacyl" refers to the group -NRC(O)R where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

5       The term "aminoacyloxy" or "alkoxycarbonylamino" refers to the group -NRC(O)OR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

10       The term "acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, aryl-C(O)O-, heteroaryl-C(O)O-, and heterocyclic-C(O)O- wherein alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, and heterocyclic are as defined herein.

15       The term "aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 20 carbon atoms having a single ring (e.g., phenyl) or multiple condensed (fused) rings, wherein at least one ring is aromatic (e.g., naphthyl, dihydrophenanthrenyl, fluorenyl, or anthryl). Preferred aryls include phenyl, naphthyl and the like.

20       Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxy, carboxyalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, aminoacyloxy, oxyacylamino, sulfonamide, thioalkoxy, substituted thioalkoxy, 25       thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-heteroaryl and trihalomethyl. Preferred aryl substituents include alkyl, alkoxy, halo, cyano, nitro,

trihalomethyl, and thioalkoxy.

The term "aryloxy" refers to the group aryl-O- wherein the aryl group is as defined above including optionally substituted aryl groups as also defined above.

5 The term "arylene" refers to the diradical derived from aryl (including substituted aryl) as defined above and is exemplified by 1,2-phenylene, 1,3-phenylene, 1,4-phenylene, 1,2-naphthylene and the like.

The term "amino" refers to the group -NH<sub>2</sub>.

10 The term "substituted amino" refers to the group -NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, cycloalkenyl, substituted cycloalkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic provided that both R's are not hydrogen.

15 "Amino acid" refers to any of the naturally occurring amino acids (e.g. Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D, L, or DL form. The side chains of naturally occurring amino acids are well known in the art and include, for example, hydrogen (e.g., as in glycine), alkyl (e.g., as in alanine, valine, leucine, isoleucine, proline), substituted alkyl (e.g., as in threonine, serine, methionine, cysteine, aspartic acid, asparagine, glutamic acid, glutamine, arginine, and lysine), alkaryl (e.g., as in phenylalanine and tryptophan),  
20 substituted arylalkyl (e.g., as in tyrosine), and heteroarylalkyl (e.g., as in histidine).

The term "carboxy" refers to -COOH.

The term "C-terminus" as it relates to a glycopeptide is well understood in the art. For example, for a glycopeptide of formula I, the C-terminus is the position substituted by the group R<sup>3</sup>.

25 The term "dicarboxy-substituted alkyl" refers to an alkyl group substituted with two carboxy groups. This term includes, by way of example, -CH<sub>2</sub>(COOH)CH<sub>2</sub>COOH and -CH<sub>2</sub>(COOH)CH<sub>2</sub>CH<sub>2</sub>COOH.



The term "carboxyalkyl" or "alkoxycarbonyl" refers to the groups  
"-C(O)O-alkyl", "-C(O)O-substituted alkyl", "-C(O)O-cycloalkyl", "-C(O)O-substituted  
cycloalkyl", "-C(O)O-alkenyl", "-C(O)O-substituted alkenyl", "-C(O)O-alkynyl" and  
"-C(O)O-substituted alkynyl" where alkyl, substituted alkyl, cycloalkyl, substituted  
5 cycloalkyl, alkenyl, substituted alkenyl, alkynyl and substituted alkynyl are as  
defined herein.

The term "cycloalkyl" refers to cyclic alkyl groups of from 3 to 20 carbon atoms  
having a single cyclic ring or multiple condensed rings. Such cycloalkyl groups include,  
by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl,  
10 cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

The term "substituted cycloalkyl" refers to cycloalkyl groups having from  
1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting  
of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl,  
substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino,  
15 aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto,  
thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy,  
thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy,  
heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-  
substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl  
20 and -SO<sub>2</sub>-heteroaryl.

The term "cycloalkylene" refers to a diradical of a cycloalkyl group.

The term "cycloalkenyl" refers to cyclic alkenyl groups of from 4 to 20 carbon  
atoms having a single cyclic ring and at least one point of internal unsaturation.  
Examples of suitable cycloalkenyl groups include, for instance, cyclobut-2-enyl,  
25 cyclopent-3-enyl, cyclooct-3-enyl and the like.

The term "substituted cycloalkenyl" refers to cycloalkenyl groups having from 1  
to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting

of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-heteroaryl.

The term "cycloalkylene" refers to a diradical of a cycloalkyl group.

The term "halo" or "halogen" refers to fluoro, chloro, bromo and iodo.

"Haloalkyl" refers to alkyl as defined herein substituted by 1-4 halo groups as defined herein, which may be the same or different. Representative haloalkyl groups include, by way of example, trifluoromethyl, 3-fluorododecyl, 12,12,12-trifluorododecyl, 2-bromooctyl, 3-bromo-6-chloroheptyl, and the like.

The term "heteroaryl" refers to an aromatic group of from 1 to 15 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within at least one ring (if there is more than one ring).

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxy, carboxyalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-heteroaryl and

trihalomethyl. Preferred aryl substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indolizinyl or benzothienyl). Preferred heteroaryls include pyridyl, pyrrolyl and furyl.

- 5           “Heteroarylalkyl” refers to (heteroaryl)alkyl- where heteroaryl and alkyl are as defined herein. Representative examples include 2-pyridylmethyl and the like.

The term “heteroaryloxy” refers to the group heteroaryl-O-.

- 10           The term “heteroarylene” refers to the diradical group derived from heteroaryl (including substituted heteroaryl), as defined above, and is exemplified by the groups 2,6-pyridylene, 2,4-pyridylene, 1,2-quinolinyne, 1,8-quinolinyne, 1,4-benzofuranylene, 2,5-pyridinyne, 2,5-indolinyne and the like.

- 15           The term "heterocycle" or "heterocyclic" refers to a monoradical saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 40 carbon atoms and from 1 to 10 hetero atoms, preferably 1 to 4 heteroatoms, selected from nitrogen, sulfur, phosphorus, and/or oxygen within the ring.

- 20           Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, oxo (O=), and -SO<sub>2</sub>-heteroaryl. Such heterocyclic groups can have a single ring or multiple condensed rings. Preferred heterocyclics include morpholino, piperidinyl, and the like.
- 25

Examples of nitrogen heterocycles and heteroaryls include, but are not limited to, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, morpholino, piperidinyl, tetrahydrofuranyl, and the like as well as N-alkoxy-nitrogen containing heterocycles.

Another class of heterocyclics is known as "crown compounds" which refers to a specific class of heterocyclic compounds having one or more repeating units of the formula  $[-(\text{CH}_2)_a\text{A}-]$  where  $a$  is equal to or greater than 2, and A at each separate occurrence can be O, N, S or P. Examples of crown compounds include, by way of example only,  $[-(\text{CH}_2)_3\text{-NH-}]_3$ ,  $[-((\text{CH}_2)_2\text{-O})_4-((\text{CH}_2)_2\text{-NH})_2]$  and the like. Typically such crown compounds can have from 4 to 10 heteroatoms and 8 to 40 carbon atoms.

The term "heterocyclene" refers to a diradical of a heterocycle group

The term "heterocyclooxy" refers to the group heterocyclic-O-.

The term "thioheterocyclooxy" refers to the group heterocyclic-S-.

The term "oxyacylamino" or "aminocarbonyloxy" refers to the group  $-\text{OC}(\text{O})\text{NRR}$  where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

The term "prodrug" is well understood in the art and includes compounds that are converted to pharmaceutically active compounds of the invention in a mammalian system. For example, see *Remington's Pharmaceutical Sciences*, **1980**, vol. 16, Mack Publishing Company, Easton, Pennsylvania, 61 and 424.

The term "saccharide group" refers to an oxidized, reduced or substituted saccharide monoradical covalently attached to the glycopeptide or other compound via

any atom of the saccharide moiety, preferably via the aglycone carbon atom. The term includes amino-containing saccharide groups. Representative saccharides include, by way of illustration, hexoses such as D-glucose, D-mannose, D-xylose, D-galactose, vancosamine, 3-desmethyl-vancosamine, 3-epi-vancosamine, 4-epi-vancosamine, 5 acosamine, actinosamine, daunosamine, 3-epi-daunosamine, ristosamine, D-glucamine, N-methyl-D-glucamine, D-glucuronic acid, *N*-acetyl-D-glucosamine, *N*-acetyl-D-galactosamine, sialic acid, iduronic acid, L-fucose, and the like; pentoses such as D-ribose or D-arabinose; ketoses such as D-ribulose or D-fructose; disaccharides such as 2-O-( $\alpha$ -L-vancosaminyl)- $\beta$ -D-glucopyranose, 2-O-(3-desmethyl- $\alpha$ -L-vancosaminyl)- $\beta$ -10 D-glucopyranose, sucrose, lactose, or maltose; derivatives such as acetals, amines, acylated, sulfated and phosphorylated sugars; oligosaccharides having from 2 to 10 saccharide units. For the purposes of this definition, these saccharides are referenced using conventional three letter nomenclature and the saccharides can be either in their open or preferably in their pyranose form.

15 The term "amino-containing saccharide group" refers to a saccharide group having an amino substituent. Representative amino-containing saccharides include L-vancosamine, 3-desmethyl-vancosamine, 3-epi-vancosamine, 4-epi-vancosamine, acosamine, actinosamine, daunosamine, 3-epi-daunosamine, ristosamine, *N*-methyl-D-glucamine and the like.

20 The term "spiro-attached cycloalkyl group" refers to a cycloalkyl group attached to another ring via one carbon atom common to both rings.

The term "stereoisomer" as it relates to a given compound is well understood in the art, and refers another compound having the same molecular formula, wherein the atoms making up the other compound differ in the way they are oriented in space, but 25 wherein the atoms in the other compound are like the atoms in the given compound with respect to which atoms are joined to which other atoms (e.g. an enantiomer, a diastereomer, or a geometric isomer). See for example, *Morrison and Boyde Organic*

*Chemistry, 1983, 4th ed.*, Allyn and Bacon, Inc., Boston, Mass., page 123

The term "sulfonamide" refers to a group of the formula  $-\text{SO}_2\text{NRR}$ , where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

5       The term "thiol" refers to the group  $-\text{SH}$ .

The term "thioalkoxy" refers to the group  $-\text{S-alkyl}$ .

The term "substituted thioalkoxy" refers to the group  $-\text{S-substituted alkyl}$ .

The term "thioaryloxy" refers to the group  $\text{aryl-S-}$  wherein the aryl group is as defined above including optionally substituted aryl groups also defined above.

10       The term "thioheteroaryloxy" refers to the group  $\text{heteroaryl-S-}$  wherein the heteroaryl group is as defined above including optionally substituted aryl groups as also defined above.

The term "thioether derivatives" when used to refer to the glycopeptide compounds of this invention includes thioethers ( $-\text{S-}$ ), sulfoxides ( $-\text{SO-}$ ) and sulfones  
15 ( $-\text{SO}_2-$ ).


As to any of the above groups which contain one or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible. In addition, the compounds of this invention include all stereochemical isomers arising from the  
20 substitution of these compounds.

"Cyclodextrin" refers to cyclic molecules containing six or more  $\alpha$ -D-glucopyranose units linked at the 1,4 positions by  $\alpha$  linkages as in amylose.  $\beta$ -Cyclodextrin or cycloheptaamylose contains seven  $\alpha$ -D-glucopyranose units. As used herein, the term "cyclodextrin" also includes cyclodextrin derivatives such as  
25 hydroxypropyl and sulfobutyl ether cyclodextrins. Such derivatives are described for example, in U.S. Patent Nos. 4,727,064 and 5,376,645. One preferred cyclodextrin is hydroxypropyl  $\beta$ -cyclodextrin having a degree of substitution of from about 4.1-5.1 as

measured by FTIR. Such a cyclodextrin is available from Cerestar (Hammond, Indiana, USA) under the name Cavitron™ 82003.

“Glycopeptide” refers to oligopeptides (e.g. heptapeptide) antibiotics, characterized by a multi-ring peptide core optionally substituted with saccharide groups, such as vancomycin. Examples of glycopeptides included in this definition may be found in “Glycopeptides Classification, Occurrence, and Discovery”, by Raymond C. Rao and Louise W. Crandall, (“Drugs and the Pharmaceutical Sciences” Volume 63, edited by Ramakrishnan Nagarajan, published by Marcel Dekker, Inc.). Additional examples of glycopeptides are disclosed in U.S. Patent Nos. 4,639,433; 4,643,987; 4,497,802; 4,698,327; 5,591,714; 5,840,684; and 5,843,889; in EP 0 802 199; EP 0 801 075; EP 0 667 353; WO 97/28812; WO 97/38702; WO 98/52589; WO 98/52592; and in *J. Amer. Chem. Soc.*, **1996**, *118*, 13107-13108; *J. Amer. Chem. Soc.*, **1997**, *119*, 12041-12047; and *J. Amer. Chem. Soc.*, **1994**, *116*, 4573-4590. Representative glycopeptides include those identified as A477, A35512, A40926, A41030, A42867, A47934, A80407, A82846, A83850, A84575, AB-65, Actaplanin, Actinoidin, Ardacin, Avoparcin, Azureomycin, Balhimycin, Chloroorientein, Chloropolysporin, Decaplanin, *N*-demethylvancomycin, Eremomycin, Galacardin, Helvecardin, Izupeptin, Kibdelin, LL-AM374, Mannopectin, MM45289, MM47756, MM47761, MM49721, MM47766, MM55260, MM55266, MM55270, MM56597, MM56598, OA-7653, Orenticin, Parvodicin, Ristocetin, Ristomycin, Synmonicin, Teicoplanin, UK-68597, UK-69542, UK-72051, Vancomycin, and the like. The term “glycopeptide” as used herein is also intended to include the general class of peptides disclosed above on which the sugar moiety is absent, i.e. the aglycone series of glycopeptides. For example, removal of the disaccharide moiety appended to the phenol on vancomycin by mild hydrolysis gives vancomycin aglycone. Also within the scope of the invention are glycopeptides that have been further appended with additional saccharide residues, especially aminoglycosides, in a manner similar to vancosamine.

"Vancomycin" refers to a glycopeptide antibiotic having the formula:



When describing vancomycin derivatives, the term "N<sup>van</sup>-" indicates that a substituent is covalently attached to the amino group of the vancosamine moiety of vancomycin.

Similarly, the term "N<sup>leu</sup>-" indicates that a substituent is covalently attached to the amino group of the leucine moiety of vancomycin.

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted" means that a group may or may not be substituted with the described substituent.

As used herein, the terms "inert organic solvent" or "inert solvent" or "inert diluent" mean a solvent or diluent which is essentially inert under the conditions of the reaction in which it is employed as a solvent or diluent. Representative examples of



materials which may be used as inert solvents or diluents include, by way of illustration, benzene, toluene, acetonitrile, tetrahydrofuran ("THF"), dimethylformamide ("DMF"), chloroform (" $\text{CHCl}_3$ "), methylene chloride (or dichloromethane or " $\text{CH}_2\text{Cl}_2$ "), diethyl ether, ethyl acetate, acetone, methylethyl ketone, methanol, ethanol, propanol, isopropanol, tert-butanol, dioxane, pyridine, and the like. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert solvents.

The term "nitrogen-linked" means a group or substituent is attached to the remainder of a compound (e.g. a compound of formula I) through a bond to a nitrogen of the group or substituent.

"Pharmaceutically acceptable salt" means those salts which retain the biological effectiveness and properties of the parent compounds and which are not biologically or otherwise harmful as the dosage administered. The compounds of this invention are capable of forming both acid and base salts by virtue of the presence of amino and carboxy groups respectively.

Pharmaceutically acceptable base addition salts may be prepared from inorganic and organic bases. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines, substituted amines including naturally-occurring substituted amines, and cyclic amines, including isopropylamine, trimethyl amine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, tromethamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, N-alkylglucamines, theobromine, purines, piperazine, piperidine, and N-ethylpiperidine. It should also be understood that other carboxylic acid derivatives would be useful in the practice of this invention, for example carboxylic acid amides, including carboxamides, lower alkyl carboxamides, di(lower alkyl) carboxamides, and the like.

Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids. Salts derived from inorganic acids include hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like. Salts derived from organic acids include acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

The compounds of this invention typically contain one or more chiral centers. Accordingly, this invention is intended to include racemic mixtures, diastereomers, enantiomers and mixture enriched in one or more stereoisomer. The scope of the invention as described and claimed encompasses the racemic forms of the compounds as well as the individual enantiomers and non-racemic mixtures thereof.

The term "treatment" as used herein includes any treatment of a condition or disease in an animal, particularly a mammal, more particularly a human, and includes:

- (i) preventing the disease or condition from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it;
- (ii) inhibiting the disease or condition, i.e. arresting its development; relieving the disease or condition, i.e. causing regression of the condition; or relieving the conditions caused by the disease, i.e. symptoms of the disease.

The term "disease state which is alleviated by treatment with a broad spectrum antibacterial" or "bacterial disease" as used herein is intended to cover all disease states which are generally acknowledged in the art to be usefully treated with a broad spectrum antibacterial in general, and those disease states which have been found to be usefully treated by the specific antibacterials of this invention. Such disease states include, but are not limited to, treatment of a mammal afflicted with pathogenic bacteria, in particular staphylococci (methicillin sensitive and resistant), streptococci (penicillin

sensitive and resistant), enterococci (vancomycin sensitive and resistant), and *Clostridium difficile*

The term "therapeutically effective amount" refers to that amount which is sufficient to effect treatment, as defined herein, when administered to a mammal in need  
5 of such treatment. The therapeutically effective amount will vary depending on the subject and disease state being treated, the severity of the affliction and the manner of administration, and may be determined routinely by one of ordinary skill in the art.

The term "protecting group" or "blocking group" refers to any group which, when bound to one or more hydroxyl, thiol, amino, carboxy or other groups of the  
10 compounds, prevents undesired reactions from occurring at these groups and which protecting group can be removed by conventional chemical or enzymatic steps to reestablish the hydroxyl, thio, amino, carboxy or other group. The particular removable blocking group employed is not critical and preferred removable hydroxyl blocking groups include conventional substituents such as allyl, benzyl, acetyl, chloroacetyl,  
15 thiobenzyl, benzylidene, phenacyl, *t*-butyl-diphenylsilyl and any other group that can be introduced chemically onto a hydroxyl functionality and later selectively removed either by chemical or enzymatic methods in mild conditions compatible with the nature of the product. Protecting groups are disclosed in more detail in T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis" 3<sup>rd</sup> Ed., 1999, John Wiley and Sons,  
20 N.Y.

Preferred removable amino blocking groups include conventional substituents such as *t*-butoxycarbonyl (*t*-BOC), benzyloxycarbonyl (CBZ), fluorenylmethoxycarbonyl (Fmoc), allyloxycarbonyl (ALOC) and the like, which can be removed by conventional conditions compatible with the nature of the product.


25 Preferred carboxy protecting groups include esters such as methyl, ethyl, propyl, *t*-butyl etc. which can be removed by mild conditions compatible with the nature of the product.

### General Synthetic Procedures


The glycopeptide compounds of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction  
5 temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

Additionally, as will be apparent to those skilled in the art, conventional  
10 protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection are well known in the art. For example, numerous protecting groups, and their  
15 introduction and removal, are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.

In the following reaction schemes, the glycopeptide compounds are depicted in a simplified form as a box "G" that shows the carboxy terminus labeled [C], the  
20 vancosamine amino terminus labeled [V], the "non-saccharide" amino terminus (leucine amine moiety) labeled [N], and optionally, the resorcinol moiety labeled [R] as follows:



By way of illustration, a glycopeptide compound, such as vancomycin, can be reductive alkylated as shown in the following reaction:



- where A represents R<sup>a</sup> minus one carbon atom and R<sup>a</sup>, R<sup>h</sup>, and W are as defined herein. This reaction is typically conducted by first contacting one equivalent of the glycopeptide, e.g., vancomycin, with an excess, preferably from 1.1 to 1.3 equivalents, of the aldehyde in the presence of an excess, preferably about 2.0 equivalents, of a tertiary amine, such as diisopropylethylamine (DIPEA) and the like. This reaction is typically conducted in an inert diluent, such as DMF or acetonitrile/water, at ambient temperature for about 0.25 to 2 hours until formation of the corresponding imine and/or hemiaminal is substantially complete. The resulting imine and/or hemiaminal is typically not isolated, but is reacted *in situ* with a metal hydride reducing agent, such as sodium cyanoborohydride or the like, to afford the corresponding amine. This reaction is preferably conducted by contacting the imine and/or hemiaminal with an excess, preferably about 3 equivalents, of trifluoroacetic acid, followed by about 1 to 1.2 equivalents of the reducing agent at ambient temperature in methanol or acetonitrile/water. The resulting alkylated product is readily purified by conventional procedures, such as precipitation and/or reverse-phase HPLC. Surprisingly, by forming the imine and/or hemiaminal in the presence of a trialkyl amine, and then acidifying

with trifluoroacetic acid before contact with the reducing agent, the selectivity for the reductive alkylation reaction is greatly improved, i.e., reductive alkylation at the amino group of the saccharide (e.g., vancosamine) is favored over reductive alkylation at the N-terminus (e.g., the leucynyl group) by at least 10:1, more preferably 20:1.

5           The above process is a significant improvement over previous methods for selectively alkylating an amino saccharide group of a glycopeptide antibiotic. Thus, the present invention also provides a method for alkylating a glycopeptide that comprises a saccharide-amine comprising:

10           combining an aldehyde or ketone, a suitable base, and the glycopeptide, to provide a reaction mixture;

          acidifying the reaction mixture; and

          combining the reaction mixture with a suitable reducing agent, to provide a glycopeptide that is alkylated at the saccharide-amine. Preferably, the glycopeptide comprises at least one amino group other than the saccharide-amine.

15           Preferably, the reductive alkylation at the saccharide-amine is favored over reductive alkylation at another amino group of the glycopeptide by at least about 10:1; and more preferably, by at least about 15:1 or about 20:1.

          The reductive alkylation process of the invention is typically carried out in the presence of a suitable solvent or combination of solvents, such as, for example, a  
20           halogenated hydrocarbon (e.g. methylene chloride), a linear or branched ether (e.g. diethyl ether, tetrahydrofuran), an aromatic hydrocarbon (e.g. benzene or toluene), an alcohol (methanol, ethanol, or isopropanol), dimethylsulfoxide (DMSO), N,N-dimethylformamide, acetonitrile, water, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, tetramethyl urea, N,N-dimethylacetamide, diethylformamide (DMF), 1-  
25           methyl-2-pyrrolidinone, tetramethylenesulfoxide, glycerol, ethyl acetate, isopropyl acetate, N,N-dimethylpropylene urea (DMPU) or dioxane. Preferably the alkylation is carried out in acetonitrile/water, or DMF/methanol.

Preferably the reduction (i.e. treatment with the reducing agent) is carried out in the presence of a protic solvent, such as, for example, an alcohol (e.g. methanol, ethanol, propanol, isopropanol, or butanol), water, or the like.

5 The reductive alkylation process of the invention can be carried out at any suitable temperature from the freezing point to the reflux temperature of the reaction mixture. Preferably the reaction is carried out at a temperature in the range of about 0 °C to about 100 °C. More preferably at a temperature in a range of about 0 °C to about 50 °C, or in a range of about 20 °C to about 30 °C.

10 Any suitable base can be employed in the reductive alkylation process of the invention. Suitable bases include tertiary amines (e.g. diisopropylethylamine, N-methylmorpholine or triethylamine) and the like.

Any suitable acid can be used to acidify the reaction mixture. Suitable acids include carboxylic acids (e.g. acetic acid, trichloroacetic acid, citric acid, formic acid, or trifluoroacetic acid), mineral acids (e.g. hydrochloric acid, sulfuric acid, or phosphoric acid), and the like. A preferred acid is trifluoroacetic acid.

15

Suitable reducing agents for carrying out reductive alkylation process of the invention are known in the art. Any suitable reducing agent can be employed in the methods of the invention, provided it is compatible with the functionality present in the glycopeptide. For example, suitable reducing agents include sodium cyanoborohydride, triacetoxyborohydride, pyridine/borane, sodium borohydride, and zinc borohydride.

20

The reduction can also be carried out in the presence of a transition metal catalyst (e.g. palladium or platinum) in the presence of a hydrogen source (e.g. hydrogen gas or cyclohexadiene). See for example, March, *Advanced Organic Chemistry*, Fourth Edition, John Wiley & Sons, New York (1992), 899-900.


25 If desired, the glycopeptide compounds of the invention can also be prepared in a step-wise manner in which a precursor to the  $-R^g-W-R^h$  group is first attached the glycopeptide by reductive alkylation, followed by subsequent elaboration of the attached

precursor using conventional reagent and procedures to form the  $-R^g-W-R^h$  group. Additionally, ketones may also be employed in the above-described reductive alkylation reactions to afford  $\alpha$ -substituted amines.

Any glycopeptide having an amino group may be employed in these reductive alkylation reactions. Such glycopeptides are well-known in the art and are either commercially available or may be isolated using conventional procedures. Suitable glycopeptides are disclosed, by way of example, in U.S. Patent Nos. 3,067,099; 3,338,786; 3,803,306; 3,928,571; 3,952,095; 4,029,769; 4,051,237; 4,064,233; 4,122,168; 4,239,751; 4,303,646; 4,322,343; 4,378,348; 4,497,802; 4,504,467; 4,542,018; 4,547,488; 4,548,925; 4,548,974; 4,552,701; 4,558,008; 4,639,433; 4,643,987; 4,661,470; 4,694,069; 4,698,327; 4,782,042; 4,914,187; 4,935,238; 4,946,941; 4,994,555; 4,996,148; 5,187,082; 5,192,742; 5,312,738; 5,451,570; 5,591,714; 5,721,208; 5,750,509; 5,840,684; and 5,843,889. Preferably, the glycopeptide employed in the above reaction is vancomycin.

As illustrated in the following scheme, an aminoalkyl sidechain at the resorcinol moiety of a glycopeptide, such as vancomycin, can be introduced via a Mannich reaction (in this scheme, the resorcinol moiety is shown for clarity). In this reaction, an amine ( $NHR^cR^c$ ) and an aldehyde ( $CH_2O$ ), such as formalin (a source of formaldehyde), are reacted with the glycopeptide under basic conditions to give the glycopeptide derivative.






Compounds of the invention comprising a sulfoxide or sulfone can be prepared from the corresponding thio compounds using conventional reagents and procedures. Suitable reagents for oxidizing a thio compound to a sulfoxide include, by way of example, hydrogen peroxide, peracides such as 3-chloroperoxybenzoic acid (MCPBA), sodium periodate, sodium chlorite, sodium hypochlorite, calcium hypochlorite, tert-butyl hypochlorite and the like. Chiral oxidizing reagents, (optically active reagents) may also be employed to provide chiral sulfoxides. Such optically active reagents are well-known in the art and include, for example, the reagents described in Kagen et al., *Synlett.*, **1990**, 643-650.

The aldehydes and ketones employed in the above reactive alkylation reactions are also well-known in the art and are either commercially available or can be prepared by conventional procedures using commercially available starting materials and conventional reagents (for example see March, *Advanced Organic Chemistry*, Fourth Edition, John Wiley & Sons, New York (1992), and references cited therein).

For example, aldehydes comprising a disulfide group are either commercially available or can be prepared by conventional procedures using commercially available starting materials and reagents. Additionally, aldehydes comprising a disulfide group can be prepared as illustrated in Scheme I.


SCHEME 1



Activation of compound 1 with diethyl azodicarboxylate (DEAD) and treatment with thiol (RSH) provides disulfide 2, which can be reduced, for example with diisobutylaluminum hydride (DIBAL-H) to provide aldehyde 3. Alternatively, intermediate compound 2 can be prepared from compound 1 by iodine oxidation in the presence of RSH.

Aldehydes comprising a thioester group are either commercially available or can be prepared by conventional procedures using commercially available starting materials and reagents. Additionally, aldehydes comprising a thioester group can be prepared as illustrated in the following Scheme 2.

SCHEME 2



- 5 Treatment of bromide 4 with thiourea provides thiol 5, which can be esterified by treatment with  $\text{RCOCl}$  to provide thioester 6. Subsequent acid catalyzed hydrolysis of the acetal provides aldehyde 9. Alternatively, ester 7 can be treated with  $\text{RCOCl}$  to provide a compound of formula 8, which can subsequently be reduced, for example using DIBAL-H, to provide the aldehyde 9.

Additional details and other methods for preparing the compounds of the invention are described in the Examples below.

### Pharmaceutical Compositions

- 10 This invention also includes pharmaceutical composition containing the novel glycopeptide compounds of this invention. Accordingly, the glycopeptide compound, preferably in the form of a pharmaceutically acceptable salt, can be formulated for oral

or parenteral administration for the therapeutic or prophylactic treatment of bacterial infections.

By way of illustration, the glycopeptide compound can be admixed with conventional pharmaceutical carriers and excipients and used in the form of tablets, capsules, elixirs, suspensions, syrups, wafers, and the like. Such pharmaceutical compositions will contain from about 0.1 to about 90% by weight of the active compound, and more generally from about 10 to about 30%. The pharmaceutical compositions may contain common carriers and excipients, such as corn starch or gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride, and alginic acid. Disintegrators commonly used in the formulations of this invention include croscarmellose, microcrystalline cellulose, corn starch, sodium starch glycolate and alginic acid.

A liquid composition will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s), for example ethanol, glycerine, sorbitol, non-aqueous solvent such as polyethylene glycol, oils or water, optionally with a suspending agent, a solubilizing agent (such as a cyclodextrin), preservative, surfactant, wetting agent, flavoring or coloring agent. Alternatively, a liquid formulation can be prepared from a reconstitutable powder.

For example a powder containing active compound, suspending agent, sucrose and a sweetener can be reconstituted with water to form a suspension; and a syrup can be prepared from a powder containing active ingredient, sucrose and a sweetener.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid compositions. Examples of such carriers include magnesium stearate, starch, lactose, sucrose, microcrystalline cellulose and binders, for example polyvinylpyrrolidone. The tablet can also be provided with a color film coating, or color included as part of the carrier(s). In

addition, active compound can be formulated in a controlled release dosage form as a tablet comprising a hydrophilic or hydrophobic matrix.

A composition in the form of a capsule can be prepared using routine encapsulation procedures, for example by incorporation of active compound and excipients into a hard gelatin capsule. Alternatively, a semi-solid matrix of active compound and high molecular weight polyethylene glycol can be prepared and filled into a hard gelatin capsule; or a solution of active compound in polyethylene glycol or a suspension in edible oil, for example liquid paraffin or fractionated coconut oil can be prepared and filled into a soft gelatin capsule.

Tablet binders that can be included are acacia, methylcellulose, sodium carboxymethylcellulose, poly-vinylpyrrolidone (Povidone), hydroxypropyl methylcellulose, sucrose, starch and ethylcellulose. Lubricants that can be used include magnesium stearate or other metallic stearates, stearic acid, silicone fluid, talc, waxes, oils and colloidal silica.

Flavoring agents such as peppermint, oil of wintergreen, cherry flavoring or the like can also be used. Additionally, it may be desirable to add a coloring agent to make the dosage form more attractive in appearance or to help identify the product.

The compounds of the invention and their pharmaceutically acceptable salts that are active when given parenterally can be formulated for intramuscular, intrathecal, or intravenous administration.

A typical composition for intra-muscular or intrathecal administration will consist of a suspension or solution of active ingredient in an oil, for example arachis oil or sesame oil. A typical composition for intravenous or intrathecal administration will consist of a sterile isotonic aqueous solution containing, for example active ingredient and dextrose or sodium chloride, or a mixture of dextrose and sodium chloride. Other examples are lactated Ringer's injection, lactated Ringer's plus dextrose injection, Normosol-M and dextrose, Isolyte E, acylated Ringer's injection, and the like.

Optionally, a co-solvent, for example, polyethylene glycol; a chelating agent, for example, ethylenediamine tetracetic acid; a solubilizing agent, for example, a cyclodextrin; and an anti-oxidant, for example, sodium metabisulphite, may be included in the formulation. Alternatively, the solution can be freeze dried and then reconstituted with a suitable solvent just prior to administration.

In a preferred embodiment, the glycopeptide derivatives of this invention are formulated in an aqueous solution containing a cyclodextrin. In another preferred embodiment the glycopeptide derivatives of this invention are formulated as a lyophilized powder containing a cyclodextrin or as a sterile powder containing a cyclodextrin. Preferably, the cyclodextrin is hydroxypropyl- $\beta$ -cyclodextrin or sulfobutyl ether  $\beta$ -cyclodextrin; more preferably, the cyclodextrin is hydroxypropyl- $\beta$ -cyclodextrin. Typically, in an injectable solution, the cyclodextrin will comprise about 1 to 25 weight percent; preferably, about 2 to 10 weight percent; more preferable, about 4 to 6 weight percent, of the formulation. Additionally, the weight ratio of the cyclodextrin to the glycopeptide derivative will preferably be from about 1:1 to about 10:1.

The compounds of the invention and their pharmaceutically acceptable salts which are active on rectal administration can be formulated as suppositories. A typical suppository formulation will generally consist of active ingredient with a binding and/or lubricating agent such as a gelatin or cocoa butter or other low melting vegetable or synthetic wax or fat.

The compounds of this invention and their pharmaceutically acceptable salts which are active on topical administration can be formulated as transdermal compositions or transdermal delivery devices ("patches"). Such compositions include, for example, a backing, active compound reservoir, a control membrane, liner and contact adhesive. Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of

pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

5 The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of  
10 the patient's symptoms, and the like.

Suitable doses are in the general range of from 0.01-100 mg/kg/day, preferably 0.1-50 mg/kg/day. For an average 70 kg human, this would amount to 0.7 mg to 7g per day, or preferably 7 mg to 3.5g per day. A more preferred dose for a human is about 500 mg to about 2g per day.

15 Other suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

The following formulation examples illustrate representative pharmaceutical compositions of the present invention.

Formulation Example A

This example illustrates the preparation of a representative pharmaceutical composition for oral administration of a compound of this invention:

	Ingredients	Quantity per tablet, (mg)
5	-----	-----
	Active Compound	200
	Lactose, spray-dried	148
	Magnesium stearate	2
	-----	-----

10 The above ingredients are mixed and introduced into a hard-shell gelatin capsule.

Formulation Example B

This example illustrates the preparation of another representative pharmaceutical composition for oral administration of a compound of this invention:

	Ingredients	Quantity per tablet, (mg)
15	-----	-----
	Active Compound	400
	Cornstarch	50
	Lactose	145
	Magnesium stearate	5
	-----	-----

20 The above ingredients are mixed intimately and pressed into single scored tablets.

Formulation Example C

This example illustrates the preparation of a representative pharmaceutical composition for oral administration of a compound of this invention.

25 An oral suspension is prepared having the following composition.



Ingredients

	Active Compound	1.0 g
	Fumaric acid	0.5 g
5	Sodium chloride	2.0 g
	Methyl paraben	0.1 g
	Granulated sugar	25.5 g
	Sorbitol (70% solution)	12.85 g
	Veegum K (Vanderbilt Co.)	1.0 g
10	Flavoring	0.035 ml
	Colorings	0.5 mg
	Distilled water	q.s. to 100 mL

Formulation Example D

15 This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

An injectable preparation buffered to a pH of 4 is prepared having the following composition:

Ingredients

20	Active Compound	0.2 g
	Sodium Acetate Buffer Solution (0.4 M)	2.0 mL
	HCl (1N)	q.s. to pH 4
	Water (distilled, sterile)	q.s. to 20 mL
25		

Formulation Example E

This example illustrates the preparation of a representative pharmaceutical composition for injection of a compound of this invention.

30 A reconstituted solution is prepared by adding 20 mL of sterile water to 1 g of the compound of this invention. Before use, the solution is then diluted with 200 mL of an intravenous fluid that is compatible with the active compound. Such fluids are

chosen from 5% dextrose solution, 0.9% sodium chloride, or a mixture of 5% dextrose and 0.9% sodium chloride. Other examples are lactated Ringer's injection, lactated Ringer's plus 5% dextrose injection, Normosol-M and 5% dextrose, Isolyte E, and acylated Ringer's injection

5 Formulation Example F

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

An injectable preparation is prepared having the following composition:

10

Ingredients	
Active Compound	0.1-5.0 g
Hydroxypropyl- $\beta$ -cyclodextrin	1-25 g
5% Aqueous Dextrose Solution (sterile)	q.s. to 100 mL

15 The above ingredients are blended and the pH is adjusted to  $3.5 \pm 0.5$  using 0.5 N HCl or 0.5 N NaOH.

Formulation Example G

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

20 A frozen solution suitable for injection is prepared having the following composition:

25

Frozen Solution	
Active Compound	250 mg to 1000 mg
Hydroxypropyl- $\beta$ -cyclodextrin	250 mg to 10 g
Excipients – e.g., dextrose	0-50 g
Water for Injection	10-100 mL

The weight ratio of hydroxypropyl- $\beta$ -cyclodextrin to the active compound will typically be from about 1:1 to about 10:1.

Representative Procedure: Hydroxypropyl- $\beta$ -cyclodextrin and excipients, if any, are dissolved in about 80% of the water for injection and the active compound is added and dissolved. The pH is adjusted with 1 M sodium hydroxide to  $4.7 \pm 0.3$  and the volume is then adjusted to 95% of the final volume with water for injection. The pH is checked and adjusted, if necessary, and the volume is adjusted to the final volume with water for injection. The formulation is then sterile filtered through a 0.22 micron filter and placed into a sterile vial under aseptic conditions. The vial is capped, labeled and stored frozen.

#### Formulation Example H

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

A lyophilized powder useful for preparing an injectable solution is prepared having the following composition:

##### Lyophilized Powder

Active Compound	250 mg to 1000 mg
Hydroxypropyl- $\beta$ -cyclodextrin	250 mg to 10 g
Excipients – e.g., mannitol, sucrose and/or lactose	0-50 g
Buffer agent – e.g., citrate	0-500 mg

The weight ratio of hydroxypropyl- $\beta$ -cyclodextrin to the active compound will typically be from about 1:1 to about 10:1.

Representative Procedure: Hydroxypropyl- $\beta$ -cyclodextrin and excipients and/or buffering agents, if any, are dissolved in about 60% of the water for injection.

The active compound is added and dissolved and the pH is adjusted with 1 M sodium hydroxide to 4.0-5.0 and the volume is adjusted to 95% of the final

5 volume with water for injection. The pH is checked and adjusted, if necessary, and the volume is adjusted to the final volume with water for injection. The formulation is then sterile filtered through a 0.22 micron filter and placed into a sterile vial under aseptic conditions. The formulation is then freeze-dried using an appropriate lyophilization cycle. The vial is capped (optionally under partial vacuum or dry nitrogen), labeled and stored at room temperature or under refrigeration.

#### Formulation Example I

10 This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

A sterile powder useful for preparing an injectable solution is prepared having the following composition:

	<u>Sterile Powder</u>	
15	Active Compound	250 mg to 1000 mg
	Hydroxypropyl- $\beta$ -cyclodextrin	250 mg to 10 g <sup>1</sup>
	Excipients	To be determined

20 The weight ratio of hydroxypropyl- $\beta$ -cyclodextrin to the active will typically be from about 1:1 to about 10:1.

Representative Procedure: Hydroxypropyl- $\beta$ -cyclodextrin and the active compound (and any excipients) are dispersed into an appropriate sterile container and the container is sealed (optionally under partial vacuum or dry nitrogen), labeled and stored at room temperature or under refrigeration.

#### 25 Administration of representative formulations H and I to a Patient

The pharmaceutical formulations described in formulation examples H and I above can be administered intravenously to a patient by the appropriate medical

personnel to treat or prevent gram-positive infections. For administration, the above formulations can be reconstituted and/or diluted with a diluent, such as 5% dextrose or sterile saline, as follows:

Representative Procedure: The lyophilized powder of formulation example H (e.g., containing 1000 mg of active compound) is reconstituted with 20 mL of sterile water and the resulting solution is further diluted with 80 mL of sterile saline in a 100 mL infusion bag. The diluted solution is then administered to the patient intravenously over 30 to 120 minutes.

#### Formulation Example J

This example illustrates the preparation of a representative pharmaceutical composition for topical application of a compound of this invention.

Ingredients	grams
Active compound	0.2-10
Span 60	2
Tween 60	2
Mineral oil	5
Petrolatum	10
Methyl paraben	0.15
Propyl paraben	0.05
BHA (butylated hydroxy anisole)	0.01
Water	q.s. to 100

All of the above ingredients, except water, are combined and heated to 60°C with stirring. A sufficient quantity of water at 60°C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. 100 g.

Formulation Example K

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

A suppository totaling 2.5 grams is prepared having the following composition:

5           Ingredients

Active Compound	500 mg
Witepsol H-15*	balance

10           (\*triglycerides of saturated vegetable fatty acid; a product of Riches-Nelson, Inc., New York, N.Y.)

Utility

The glycopeptide compounds of this invention, and their pharmaceutically acceptable salts, are useful in medical treatments and exhibit biological activity, including antibacterial activity, which can be demonstrated in using the tests described herein. Such tests are well known to those skilled in the art, and are referenced and described in Lorian "Antibiotics in Laboratory Medicine", Fourth Edition, Williams and Wilkins (1991).

Accordingly, this invention provides methods for treating bacterial or infectious diseases, especially those caused by Gram-positive microorganisms, in animals. The compounds of this invention are particularly useful in treating infections caused by methicillin-resistant staphylococci. Also, the compounds are useful in treating infection due to enterococci, including vancomycin-resistant enterococci (VRE). Examples of such diseases include severe staphylococcal infections, such as staphylococcal endocarditis and staphylococcal septicemia. The animal treated may be either susceptible to, or infected with, the microorganism. The method of treatment typically comprises administering to the animal an amount of a compound of this invention which is effective for this purpose.

In practicing this method, the antibiotic can be administered in a single daily dose or in multiple doses per day. The treatment regimen may require administration over extended periods of time, for example, for several days or for from one to six weeks. The amount per administered dose or the total amount administered will depend  
5 on such factors as the nature and severity of the infection, the age and general health of the patient, the tolerance of the patient to the antibiotic and the microorganism or microorganisms in the infection.

Among other properties, the glycopeptide compounds of the invention have been found to have reduced mammalian toxicity when administered to a mammal. For  
10 example, the disulfide or thioester derivatives of the invention have been found to have reduced liver and/or kidney accumulation compared to the corresponding compounds that lack the disulfide or thioester group(s). Moreover, certain compounds of the invention are expected to have reduced nephrotoxicity. Additionally, it has been discovered that the addition of a cyclodextrin compound to a pharmaceutical  
15 composition containing the glycopeptide compounds of this invention further reduces the nephrotoxicity and/or tissue accumulation of the glycopeptide compound when administered to a mammal.

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention.

## 20 EXAMPLES

In the examples below, the following abbreviations have the following meanings. Any abbreviations not defined have their generally accepted meaning. Unless otherwise stated, all temperatures are in degrees Celsius.

ACN	=	acetonitrile
BOC, Boc	=	<i>tert</i> -butoxycarbonyl
DIBAL-H	=	diisobutylaluminum hydride

15

20

## 25

To a mixture of vancomycin (1 eq.) and the desired aldehyde (1.3 eq.) in DMF was added DIPEA (2 eq.). The reaction was stirred at ambient temperature for 1-2 hours and monitored by reverse-phase HPLC. Methanol and NaCNBH<sub>3</sub> (1 eq.) were added to the solution, followed by TFA (3 eq.). Stirring was continued for an additional hour at ambient temperature. After the reaction was complete, the methanol was removed *in vacuo*. The residue was precipitated in acetonitrile. Filtration gave the crude product which was then purified by reverse-phase HPLC. If desired, other glycopeptides antibiotics may be used in this procedure.



### General Procedure B

#### Preparation of $\text{MeOC(O)CH}_2\text{SSC}_8\text{H}_{17}$

In a  $\text{N}_2$  purged flask there was dissolved Diethyl azodicarboxylate (5.0g, 28.71mmol) in dichloromethane (150mL). A solution of Methyl thioglycolate (2.89g, 27.27mmol) in dichloromethane (50mL) was then added dropwise at room temperature over 20 minutes. After 1.5 hours stirring, a solution of octanethiol (4.8mL, 27.27mmol) in dichloromethane (50 mL) was added dropwise over 20 minutes, and the resulting solution was stirred for an additional 12 hours at room temperature. The solvent was then removed under reduced pressure, and the resulting oil was then purified over flash silica with 5% ethyl acetate in hexanes to yield  $\text{MeOC(O)CH}_2\text{SSC}_8\text{H}_{17}$  as a clear, colorless oil.

### General Procedure C

#### Preparation of $\text{OHCCH}_2\text{SSC}_8\text{H}_{17}$

$\text{MeOC(O)CH}_2\text{SSC}_8\text{H}_{17}$  (1.85g, 7.39mmol) was dissolved in dry diethyl ether (40mL) in a  $\text{N}_2$  purged flask. The solution was cooled to  $-78^\circ\text{C}$ , and DIBAL-H (8.87mL, 1.0 M in cyclohexane, 8.87mmol) was added to the cold solution dropwise via syringe. The reaction was stirred at  $-78^\circ\text{C}$  for 1h 45min, after which  $\text{H}_2\text{O}$  (5mL) was added, the cold bath removed and the solution was allowed to slowly warm to room temperature. Additional diethyl ether (60mL) was added, and the organic phase was washed with 1N HCl (3 x 100mL), dried over  $\text{MgSO}_4$ , and concentrated to yield  $\text{HC(O)CH}_2\text{SSC}_8\text{H}_{17}$  (1.58g, 97.02%) as a milky oil.

### Example 1

#### Preparation of Compound 1

Using general reductive alkylation procedure A described above, and the following amounts of materials, the title compound was prepared.

5     Amounts:

HC(O)CH<sub>2</sub>SSC<sub>8</sub>H<sub>17</sub> (1.58g, 7.17mmol)

Vancomycin (10.65g, 7.17mmol)

Diisopropylethyl amine (2.5mL, 14.20mmol)

TFA (1.66mL, 21.51mmol)

10     NaCNBH<sub>3</sub> (1.35g, 21.51mmol)

Once reduction of the imine was complete as evidenced by HPLC, the reaction mixture was poured into diethyl ether, and the resulting ppt was filtered. The crude product was purified by reverse phase HPLC to yield the title compound. MS calculated (M<sup>+</sup>) 1653.7; found (MH<sup>+</sup>)1654.4.

15     Using the above procedures and the appropriate starting materials the compounds shown in Table I were prepared. The mass spectral data for these compounds were as follows:

Compound No.	MW (freebase)	Observed MH <sup>+</sup>
1	1653.7	1654.4
2	1667.71	1668.8
3	1667.71	1668.5
4	1681.74	1682.8

## Example 2

### Determination of Antibacterial Activity

#### A. In Vitro Determination of Antibacterial Activity

##### 1. Determination of Minimal Inhibitory Concentrations (MICs)

5 Bacterial strains were obtained from either American Type Tissue Culture Collection (ATCC), Stanford University Hospital (SU), Kaiser Permanente Regional Laboratory in Berkeley (KPB), Massachusetts General Hospital (MGH), the Centers for Disease Control (CDC), the San Francisco Veterans' Administration Hospital (SFVA) or the University of California San Francisco Hospital (UCSF). Vancomycin resistant  
10 enterococci were phenotyped as Van A or Van B based on their sensitivity to teicoplanin. Some vancomycin resistant enterococci that had been genotyped as Van A, Van B, Van C1 or Van C2 were obtained from the Mayo Clinic.

Minimal inhibitory concentrations (MICs) were measured in a microdilution  
15 broth procedure under NCCLS guidelines. Routinely, the compounds were serially diluted into Mueller-Hinton broth in 96-well microtiter plates. Overnight cultures of bacterial strains were diluted based on absorbance at 600 nm so that the final concentration in each well was  $5 \times 10^5$  cfu/mL. Plates were returned to a 35°C incubator. The following day (or 24 hours in the case of Enterococci strains), MICs  
20 were determined by visual inspection of the plates. Strains routinely tested in the initial screen included methicillin-sensitive *Staphylococcus aureus* (MSSA), methicillin-resistant *Staphylococcus aureus*, methicillin-sensitive *Staphylococcus epidermidis* (MSSE), methicillin-resistant *Staphylococcus epidermidis* (MRSE), vancomycin sensitive *Enterococcus faecium* (VSE Fm), vancomycin sensitive *Enterococcus faecalis* (VSE Fs), vancomycin resistant *Enterococcus faecium* also resistant to teicoplanin  
25 (VRE Fm Van A), vancomycin resistant *Enterococcus faecium* sensitive to teicoplanin

(VRE Fm Van B), vancomycin resistant *Enterococcus faecalis* also resistant to teicoplanin (VRE Fs Van A), vancomycin resistant *Enterococcus faecalis* sensitive to teicoplanin (VRE Fs Van B), enterococcus gallinarum of the Van A genotype (VRE Gm Van A), enterococcus gallinarum of the Van C-1 genotype (VRE Gm Van C-1),  
5 enterococcus casseliflavus of the Van C-2 genotype (VRE Cs Van C-2), enterococcus flavescens of the Van C-2 genotype (VRE Fv Van C-2), and penicillin-sensitive *Streptococcus pneumoniae* (PSSP) and penicillin-resistant *Streptococcus pneumoniae* (PSRP). Because of the inability of PSSP and PSRP to grow well in Mueller-Hinton broth, MICs with those strains were determined using either TSA broth supplemented  
10 with defibrinated blood or blood agar plates. Compounds which had significant activity against the strains mentioned above were then tested for MIC values in a larger panel of clinical isolates including the species listed above as well as non-speciated coagulase negative *Staphylococcus* both sensitive and resistant to methicillin (MS-CNS and MR-CNS). In addition, they were tested for MICs against gram negative organisms, such as  
15 *Escherichia coli* and *Pseudomonas aeruginosa*.

## 2. Determination of Kill Time

Experiments to determine the time required to kill the bacteria were conducted as described in Lorian, "Antibiotics in Laboratory Medicine", Fourth Edition, Williams and Wilkins (1991). These experiments were conducted normally with both  
20 staphylococcus and enterococcus strains.

Briefly, several colonies were selected from an agar plate and grown at 35°C under constant agitation until it achieved a turbidity of approximately 1.5 and 10<sup>8</sup> CFU/mL. The sample was then diluted to about 6 x 10<sup>6</sup> CFU/mL and incubated at 35°C under constant agitation was continued. At various times aliquots were removed and  
25 five ten-fold serial dilutions were performed. The pour plate method was used to determine the number of colony forming units (CFUs).

In general, the compounds of the invention were active in the above tests *in vitro* tests and demonstrated a broad spectrum of activity.

B. In Vivo Determination of Antibacterial Activity

1. Acute Tolerability Studies in Mice

5 In these studies, a compound of this invention was administered either intravenously or subcutaneously and observed for 5-15 minutes. If there were no adverse effects, the dose was increased in a second group of mice. This dose incrementation continued until mortality occurred, or the dose was maximized. Generally, dosing began at 20 mg/kg and increased by 20 mg/kg each time until the  
10 maximum tolerated dose (MTD) is achieved.

2. Bioavailability Studies in Mice

Mice were administered a compound of this invention either intravenously or subcutaneously at a therapeutic dose (in general, approximately 50 mg/kg). Groups of animals were placed in metabolic cages so that urine and feces could be collected for  
15 analysis. Groups of animals (n = 3) were sacrificed at various times (10 min, 1 hour and 4 hours). Blood was collected by cardiac puncture and the following organs were harvested—lung, liver, heart, brain, kidney, and spleen. Tissues were weighed and prepared for HPLC analysis. HPLC analysis on the tissue homogenates and fluids was used to determine the concentration of the test compound or IiI present. Metabolic  
20 products resulting from changes to the test compound were also determined at this juncture.

3. Mouse Septicemia Model

In this model, an appropriately virulent strain of bacteria (most commonly *S. aureus*, or *E. Faecalis* or *E. Faecium*) was administered to mice (N=5 to 10 mice per

group) intraperitoneally. The bacteria was combined with hog gastric mucin to enhance virulence. The dose of bacteria (normally  $10^5$ - $10^7$ ) was that sufficient to induce mortality in all of the mice over a three day period. One hour after the bacteria was administered, a compound of this invention was administered in a single dose either IV or subcutaneously. Each dose was administered to groups of 5 to 10 mice, at doses that typically ranged from a maximum of about 20 mg/kg to a minimum of less than 1 mg/kg. A positive control (normally vancomycin with vancomycin sensitive strains) was administered in each experiment. The dose at which approximately 50% of the animals are saved was calculated from the results.

#### 4. Neutropenic Thigh Model

In this model, antibacterial activity of a compound of this invention was evaluated against an appropriately virulent strain of bacteria (most commonly *S. aureus*, or *E. Faecalis* or *E. Faecium*, sensitive or resistant to vancomycin). Mice were initially rendered neutropenic by administration of cyclophosphamide at 200 mg/kg on day 0 and day 2. On day 4 they were infected in the left anterior thigh by an IM injection of a single dose of bacteria. The mice were then administered the test compound one hour after the bacteria and at various later times (normally 1, 2.5, 4 and 24 hours) the mice were sacrificed (3 per time point) and the thigh excised, homogenized and the number of CFUs (colony forming units) were determined by plating. Blood was also plated to determine the CFUs in the blood.

#### 5. Pharmacokinetic Studies

The rate at which a compound of this invention is removed from the blood can be determined in either rats or mice. In rats, the test animals were cannulated in the jugular vein. The test compound was administered via tail vein injection, and at various time points (normally 5, 15, 30, 60 minutes and 2, 4, 6 and 24 hours) blood was

withdrawn from the cannula. In mice, the test compound was also administered via tail vein injection, and at various time points. Blood was normally obtained by cardiac puncture. The concentration of the remaining test compound was determined by HPLC.

In general, the compounds of the invention were active in the above test in vivo  
5 and demonstrated a broad spectrum of activity.

### Example 3

#### Determination of Tissue Accumulation

##### A. Tissue Distribution Using Radiolabeled Compound

This procedure is used to examine the tissue distribution, excretion and  
10 metabolism of a radiolabeled test compound in both male and female rats following intravenous infusion at 10 mg/kg. Male and female Sprague-Dawley rats (n = 2 per sex per compound) are dosed with <sup>3</sup>H-labeled test compound at 10 (400 µCi/kg) and 12.5 mg/kg (100 µCi/kg), respectively, via intravenous infusion (~2 min). The test compound is formulated in 5% hydroxypropyl-β-cyclodextrin as 2.5 mg/mL solution.  
15 Urine and feces are cage collected over 24 hours period. At 24 hours after dosing, animals are sacrificed and tissues are removed. Serum, urine and tissues are analyzed for total radioactivity by oxidation followed by liquid scintillation counting. Urine and selected tissues samples are extracted and analyzed by reverse phase HPLC with radioactive flow detector for the presence of potential metabolites.

##### 20 B. Tissue Accumulation Following Single Dose

This procedure is used to evaluate tissue distribution of a test compound in rats following single dose administration by infusion. Male Sprague-Dawley rats (n = 3 per dose groups) are dosed with 50 mg/kg of a test compound. Two formulations are used: 30% PEG 400 and 10% sulfobutylether-β-cyclodextrin. Urine samples are cage  
25 collected over 24 hours. Blood samples are collected for serum chemistry and

concentration determination. Liver and kidneys are removed for histology evaluation. One kidney and part of the liver are homogenized for concentration analysis using reverse phase HPLC with UV detection. Drug concentrations in urine and serum samples are determined by LC-MS analysis.

5                   C.       Tissue Distribution Following Multiple Doses

                  This procedure is used too evaluate the potential tissue accumulation of a test compound in rats following multiple dose administration by intravenous infusion. Male and female Sprague-Dawley rats (n = 4 per sex per dose group) are dosed with a test compound at 12.5, 25 and 50 mg/kg per day for seven days. Animals are sacrificed at  
10           day 1 (n = 3 per sex per dose group) following the last dose administered. One animal per sex per dose group is retained as recovery animal and sacrificed at day 7 following the last dose administered. The test compound is formulated in 5% hydroxypropyl- $\beta$ -cyclodextrin or 1% sucrose/4.5% dextrose. Urine samples are cage collected at days 1 and 7 post-dose. Blood samples are collected for serum chemistry and concentration  
15           determination. Liver and kidneys are removed for histology evaluation. One kidney and part of the liver are homogenized for concentration analysis using reverse phase HPLC with UV detection. Drug concentrations in urine and serum samples are determined by LC-MS analysis.

                  While the present invention has been described with reference to the specific  
20           embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications  
25           are intended to be within the scope of the claims appended hereto. Additionally, all



publications, patents, and patent documents cited hereinabove are incorporated by reference herein in full, as though individually incorporated by reference.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2